

COMPARATIVE ANALYSIS OF HOLSTEIN, BLACK-MOTLEY, ANGLER, SIMMENTAL BULLS SEMEN

**Mussabekov A. T.¹, Borovikov S. N.¹, Suranshiyev Zh. A.¹,
Shamshidin A. S.²**

¹ – S. Seifullin Kazakh Agro Technical University
Astana, Akmola Region, Republic of Kazakhstan¹

² – Republican livestock breeding centre JSC «Assyl tulik»
Akmola Region, Republic of Kazakhstan²

Obtaining semen from manufacturers is the most important technological element in artificial insemination. There are some requirements in the methods of the obtaining semen : ensuring the maximum volume of ejaculate and protecting sperm from injuries, guaranteed high sanitary quality of sperm, safety, health and reproductive capacity of males. The effectiveness of artificial insemination of females depends on many factors: the quality and quantity of sperm cells in the dose, timing and frequency of insemination, taking into account the survival of sperm and egg cells, sperm injection method, and the state of the female reproductive organs. It is important to know exactly survival time of sperm and the egg in the female reproductive system, as well as the speed and the time of sperm travel from the injection site to the egg cells, as this depends on the time and frequency of insemination.

The study was conducted in the Akmola region, Republican livestock breeding centre JSC «Assyl tulik». The purpose of the work is a comparative evaluation of bulls' sperm of different genotypes. Four experimental groups formed. In the I group sires of Holstein red-motley breed, in the II – Black-motley, in the III – Angler breed in the IV – Simmental. All conditions and feeding, mode of use were the same for all sires. Sperm was taken twice a week with doublet nozzle on substituted animals.

Sperm was investigated by conventional methods: the volume was measured with a graduated cylinder; activity – under a microscope at a magnification of 180-200 times by a 10-point scale; the concentration of sperm in 1 ml of semen was measured by using photo colorimeter FEC-M. The resulting semen was diluted by lactose-yolk diluent with glycerol and calculated the content of not less than 50 million sperm cells in a dose. After that they were frozen. Prior to packaging and after a month of storage control checking of stability of sperm to the deep cooling was made.

It is known that the results of artificial insemination are largely dependent on the quality of sperm. Therefore, an objective assessment, and thorough research of the seed is an important condition for the effectiveness of

insemination (Table). By the main breeding grounds and the degree of inheritance of economically useful properties it was found that the reproductive function of the bulls is largely determined by large genetic diversity.

Parameters	Group			
	I	II	III	IV
Amount of ejaculates for half a year	151,7±2,18	137,7±5,21	110,3±2,72	146,7±1,85
Received sperm, ml	1016,3±65,93	797,3±57,78	494,7±85,68	1010,7±78,97
The volume of ejaculate, ml	6,7±0,35	5,8±0,51	4,4±0,77	6,54±0,47
The concentration, l rd/ml	0,98±0,01	0,89±0,03	0,9±0,05	0,94±0,01
Amount of frozen semen, ml	22895,0±18,95	16613,3±11,59	10363,3±22,56	22250,0±16,31
Amount of defective sperm, ml	2,7±2,18	22,3±14,51	22,7±13,67	8,3±1,20

An important indicator of sperm in its assessing is the volume of ejaculate, which means the amount of sperm, selected by the manufacturer in one of implementations. Thus, sires of red steppe breed in the amount of ejaculates surpassed peers of black-motley breed for 14 (10.2%), Angler – 41.4 (37.5%) Simmental 5 (3.4%). The lowest values of the studied index characterized sires of Angler breed. This in turn has led to the differences in the amount of produced sperm. The lowest values of the studied index characterized sires of the III group. They yielded to the analogs of the I group to 521.6 ml, II to 302.6 ml, IV to 516 ml. Sires of Simmental breed in turn slightly inferior to the sires of the red steppe breed to 5.6 ml, but superior to the sires of the black-motley breed to 213.4 ml.

The study also found a definite effect of genotype and origin of sires on the volume of ejaculate. Thus, bulls of Holstein red-motley breed are characterized by the highest volume of ejaculate. They are slightly superior bulls of Simmental breed – 0.2 ml (3.1%). The superiority of the bulls of black-motley breed was 0.9 ml (15.55), Angler – 2.3 ml (52.3%). At the same time the bulls of black-motley breed superior bulls of Angler breed – 1.4 mL (31.8), but conceded bulls of Simmental breed 0.7 ml (12.1%). The lowest rates of volume of ejaculate are characterized sires of Angler breed.

It should be noted that the concentration of sperm in 1 ml of sires of all test groups were at the same level and amounted to 0,92-0,95 billion/ml. In the obtaining of the sperm one of important things are quality indicators of sperm and sperm viability. Thus, it was found that the amount of the highest culled-term sperm were at sires of the III group. They were superior to their peers of the II group to 0.4 ml (1,8%), the I group to 20 ml (88,1%), the VI group to 14.4 ml (63.4%). Bulls of Simmental breed, in turn, had an advantage over the bulls of red steppe breed to 5.6 ml. Bulls of Holstein red-

motley breed are characterized by the lowest number of culled sperm. As a consequence of it from the I experienced group of animals was frozen to 645 ml (2.9%) more sperm than the bulls from the IV group. Superior to the bulls of the II group was 6281.7 ml (37,8%), III – 12531,7 ml (120.9%). Bulls of the Angler breed are characterized by the lowest value of the studied index. They yielded their analogs of the II group to 6250 ml (60,3%), IV group – 11886.7 (53.4%). Sires of the Simmental breed in turn superior sires of the black-motley breed by the amount of frozen semen to 5636.7 ml (25.3%). Indicators of sperm.

Summarizing the above, we can note that the sperm parameters are directly dependent on the breed, the individual characteristics of sires. Higher quantitative and qualitative indicators has sires of Holstein and Black-motley breed. They were slightly inferior bulls of Simmental breed. The lowest sperm indexes have sires of Angler breed.