

2. - « -», 2010, . -  
 .. 69-70.
3. / ..  
 -2012, - .. 64-65.  
 606:579.67

## LACTOCOCCUS

« . » . . »

(RAPD) [1],  
 (Rep- ) [2, 3] (ITS- ) [4].

: ITS-RFLP, Rep- Lactococcus , RAPD.

Chelex 100 ( , ).

TREECON for Windows (version 1.3b).  
 16S–23S  
 Lactococcus lactis  
 650 . . .  
 Taq I Alu I ITS-RFLP  
 Rep-  
 14–25

ERIC2-1.  
 RAPD– 18 ( P16) 23 ( P15) 4  
 : 2067M-AD, 17M-AD, 705M-ADG ( A); 942MA-D, 1335 M-ADG ( B); 1940 MAD-G, 1945 MAD-G, 1947 MAD-G ( ); 1265M-A, 454M-A, 973M-A, 1240M-A ( D).  
 Lactococcus lactis  
 Rep- RAPD- (ITS-RFLP)

1. Application of RAPD analysis for identification of *Lactococcus lactis* subsp. *cremoris* strains isolated from artisanal cultures / D. Samarzija [et al.] // *Microbiol Res.* – 2002. – Vol. 157. – P.13–17.
2. Repetitive extragenic palindromic sequences: a major component of the bacterial genome / M.J. Stern [et al.] // *Cell.* – 1984. – Vol. 37. – P. 1015–1026.
3. Novel BOX repeat PCR assay for high-resolution typing of *Streptococcus pneumoniae* strains / A. Van Belkum [et al.] // *J. Clin. Microbiol.* – 1996. – V. 34. – P. 1176–1179.


4. Olive, D.M. Principles and applications of methods for DNA-based typing of microbial organisms / D.M. Olive, P. Bean // J. Clin. Microbiol. – 1999. – V. 37. – P. 1661–1669.

606:579.67

«...»

[1],

(...1).



1 - , 2 - , 3 -

1 -

m,

$P_n$ ,

(...2, )

$$F_t \geq (fP_{n1} + fP_{n2})$$

$P_{n1}$   $P_{n2}$

$$F_t \geq 2fP_n$$

$P_n$

[2],